- 13. The molecule of claim 1 wherein said molecule is an inhibitor of a mammalian BRAG2.
- **14**. An in vitro or in cellulo method for inhibiting a mammalian BRAG2, said method comprising contacting the mammalian BRAG2 with a BRAG2 inhibitor according to claim **7**.
- 15. A pharmaceutical composition comprising at least one molecule according to claim 1, said composition comprising one or more excipients and optionally one other pharmaceutically active ingredient.

**16-19**. (canceled)

- 20. A method of therapeutic treatment, said method comprising administering to a mammal in need thereof an effective amount of at least one molecule according to claim 1.
- **21**. The method of claim **20**, wherein said method is for treating a disease presenting a deregulated expression of BRAG2.
- 22. The method according to claim 20, wherein said method is for treating a disease is selected from the group consisting of a cancer, angiogenesis, diabetic retinopathy, or non-syndromic intellectual disability.
- 23. The method of claim 20, wherein said method is for treating a disease selected from the group consisting of a cancer, in particular an invasive cancer, a cancer with metastasis, a cancer resistant to an EGFR and/or ErbB2 modulator, angiogenesis, diabetic retinopathy, non-syndromic intellectual disability.
- 24. The method of claim 20, wherein said molecule is an inhibitor having one or more protein-membrane interactions and inhibiting a mammal BRAG2, and wherein said method is for the treatment of a cancer.

25. The molecule according to claim 3, wherein the alkyl is methyl (Me) or ethyl (Et); and/or the O alkyl is OMe or OFt, and/or

the O-alkyl is OMe or OEt; and/or the alkyne, is —CCH; and/or

the O-alkyne is —OCH2-CCH.

26. The molecule according to claim 4, wherein the alkyl is methyl (Me) or ethyl (Et); and/or

the O-alkyl, is OMe or OEt; and/or

the alkyne is —CCH; and/or

the O-alkyne is —OCH2-CCH.